arXiv:1404.7303v1 [math.GR] 29 Apr 2014

NEW EXAMPLES OF MIXED BEAUVILLE GROUPS

BEN FAIRBAIRN, EMILIO PIERRO

Abstract. We generalise a construction of mixed Beauville groups first given by Bauer, Catanese and Grunewald. We go on to give several examples of infinite families of characteristically simple groups that satisfy the hypotheses of our theorem and thus provide a wealth of new examples of mixed Beauville groups.

1. Introduction

We recall the following definition first made by Bauer, Catanese and Grunewald in [5, Definition 4.1].

Definition 1. *Let G be a finite group and for $x, y \in G$ define*

$$\Sigma(x,y) = \bigcup_{i=1}^{|G|} \bigcup_{g \in G} \{(x^i)^g, (y^i)^g, ((xy)^i)^g\}.$$

*A **mixed Beauville quadruple** for G is a quadruple $(G^0; a, c; g)$ consisting of a subgroup G^0 of index 2 in G; of elements $a, c \in G^0$ and of an element $g \in G$ such that*

M1 *G^0 is generated by a and c;*
M2 *$g \notin G^0$;*
M3 *for every $\gamma \in G^0$ we have that $(g\gamma)^2 \notin \Sigma(a,c)$ and*
M4 *$\Sigma(a,c) \cap \Sigma(a^g, c^g) = \{e\}$.*

*If the g is clear then we will omit this simply writing $(G^0; a, c)$ instead. If G has a mixed Beauville quadruple we say that G is a **mixed Beauville group** and call $(G^0; a, c)$ a **mixed Beauville structure** on G. Finally, the **type** of this structure is $(o(a_1), o(c_1), o(a_1c_1); o(a_2), o(c_2), o(a_2c_2))$.*

We will not discuss the corresponding 'unmixed' Beauville groups here. Beauville groups were originally introduced in connection with a class of complex surfaces of general type, known as Beauville surfaces. These surfaces possess many useful geometric properties: their automorphism groups [23] and fundamental groups [10] are relatively easy to compute and these surfaces are rigid in the sense of admitting no non-trivial deformations [6] and thus correspond to isolated points in the moduli space of surfaces of general type. Early motivation came from providing cheap counterexamples to the so-called 'Friedman-Morgan speculation' [15] but they also provide a wide class of surfaces that are unusually easy to deal with to test conjectures and provide counterexamples. A number of excellent surveys on these and closely related matters have appeared in recent years - see any of [4, 8, 9, 13, 24, 35] and the references therein.

To construct examples of mixed Beauville groups, Bauer, Catanese and Grunewald proved the following in [5, Lemma 4.5].

Theorem 2. *Let H be a perfect finite group. Let $a_1, c_1, a_2, c_2 \in H$ and define $\nu(a_i, c_i) = o(a_i)o(c_i)o(a_ic_i)$ for $i = 1, 2$. Assume that*

(1) *$o(a_1)$ and $o(c_1)$ are even;*
(2) *$\langle a_1, c_1 \rangle = H$;*
(3) *$\langle a_2, c_2 \rangle = H$;*
(4) *$\nu(a_1, c_1)$ is coprime to $\nu(a_2, c_2)$.*

Set $G := (H \times H) : \langle g \rangle$ where g is an element of order 4 that acts by interchanging the two factors; $G^0 = H \times H \times \langle g^2 \rangle$; $a := (a_1, a_2, g^2)$ and $c := (c_1, c_2, g^2)$. Then $(G^0; a, c)$ is a mixed Beauville structure on G.

The only examples of groups satisfying the hypotheses of Theorem 2 given by Bauer, Catanese and Grunewald were the alternating groups A_n 'for large n' (proved using the heavy duty machinery first employed to verify Higman's conjecture concerning alternating groups as images of triangle groups) and

the groups $SL_2(p)$ with $p \neq 2, 3, 5, 17$ prime (though their argument also does not apply to the case $p = 7$).

More generally, mixed Beauville groups have proved extremely difficult to construct. Bauer, Catanese and Grunewald show [7] that there are two groups of order 2^8 which admit a mixed Beauville structure but no group of smaller order, however, the method they use is that of checking every group computationally. Indeed, any p-group admitting a mixed Beauville structure must be a 2-group. Barker, Bosten, Peyerimhoff and Vdovina construct five new examples of mixed Beauville 2-groups in [2] and an infinite family in [3] and the aforementioned examples account for all presently known mixed Beauville groups. Non-examples of mixed Beauville groups, however, are in abundance. The following result is due to Fuertes and González-Diez [17, Lemma 5].

Lemma 3. *Let $(C \times C)/G$ be a Beauville surface of mixed type and G^0 the subgroup of G consisting of the elements which preserve each of the factors, then the order of any element $f \in G \setminus G^0$ is divisible by 4.*

Fuertes and González-Diez used the above to prove that no symmetric group is a mixed Beauville group. It is easy to see, however that this result actually rules out most almost simple groups [12] (though as the groups $P\Sigma L_2(p^2)$ with p prime show not quite all of them). Bauer, Catanese and Grunewald also show in [5, Theorem 4.3] that G^0 must be non-abelian.

In light of the above we make the following definition.

Definition 4. *Let G be a group satisfying the hypotheses of Theorem 2. Then we say that G is a **mixable Beauville group**. More specifically, G is a mixable Beauville group if G is a perfect group and there exist $x_1, y_1, x_2, y_2 \in G$ such that $o(x_1)$ and $o(y_1)$ are even; $\langle x_1, y_1 \rangle = \langle x_2, y_2 \rangle = G$ and $\nu(x_1, y_1)$ is coprime to $\nu(x_2, y_2)$. We call (x_1, y_1, x_2, y_2) a **mixable Beauville structure** for G of **type** $(o(x_1), o(y_1), o(x_1y_1); o(x_2), o(y_2), o(x_2y_2))$.*

Thus any mixable Beauville group automatically gives us a mixed Beauville group by Theorem 2. In Section 2 we will prove a generalisation of Theorem 2 that shows a given mixable Beaville group actually provides us with a wealth of mixed Beauville groups.

We remark that finding 2-generated non-mixable Beauville groups is not difficult. For example, no symmetric group is mixable (every generating set must contain elements of even order, so we cannot satisfy the conditions in Definition 4); p-groups are not mixable (to have an index 2 subgroup we must have $p = 2$ and then again the conditions in Definition 4 cannot be satisfied) and even among the finite simple groups, $PSL_2(2^n)$ for $n \geq 2$ are not mixable (the only elements of even order have order 2 and thus condition (2) of Theorem 2 cannot be satisfied). Despite this, we prove the following.

Theorem 5. *If G belongs to any of the following families of simple groups*

- *the alternating groups A_n, $n \geq 6$;*
- *the linear groups $PSL_2(q)$ with $q \geq 7$ odd;*
- *the unitary groups $U_3(q)$ with $q \geq 3$;*
- *the Suzuki groups ${}^2B_2(2^{2n+1})$ with $n \geq 1$;*
- *the small Ree groups ${}^2G_2(3^{2n+1})$ with $n \geq 1$;*
- *the large Ree groups ${}^2F_4(2^{2n+1})$ with $n \geq 1$;*
- *the Steinberg triality group ${}^3D_4(q)$ with $q \geq 2$, or;*
- *the sporadic groups (including the Tits group ${}^2F_4(2)'$),*

then G is a mixable Beauville group. Furthermore, if G is any of the above or $PSL_2(2^n)$ with $n \geq 3$ then $G \times G$ is also a mixable Beauville group.

In light of the above Theorem we make the following conjecture.

Conjecture 6. If G is a finite simple group not isomorphic to $PSL_2(2^n)$ for $n \geq 2$ then G is a mixable Beauville group.

Groups of the form G^n where G is a finite simple group are called characteristically simple groups. The study of characteristically simple Beauville groups has recently been initiated by Jones in [25, 26]. It is easy to show that the characteristically simple group $PSL_2(7) \times PSL_2(7) \times PSL_2(7)$ is not a mixable Beauville group. We thus ask the following natural question.

Question 7. If G is a simple group then for which n is G^n a mixable Beauville group?

Throughout we use the standard 'Atlas' notation for finite groups and related concepts as described in [11]. In Section 2 we discuss a generalization of Theorem 2 which enables mixable Beauville groups to define several mixed Beauville groups before turning our attention in the remaining Sections to the proof of Theorem 5.

2. Generalisation and auxiliary results

We begin with the following definition which will be key to the generating structures we demonstrate.

Definition 8. *Let G be a finite group and $x, y, z \in G$. A **hyperbolic generating triple** for G is a triple $(x, y, z) \in G \times G \times G$ such that*

1. $\frac{1}{o(x)} + \frac{1}{o(y)} + \frac{1}{o(z)} < 1$,
2. $\langle x, y, z\rangle = G$, *and;*
3. $xyz = 1$.

*The **type** of a hyperbolic generating triple (x, y, z) is the triple $(o(x), o(y), o(z))$. If at least two of $o(x), o(y)$ and $o(z)$ are even then we call (x, y, z) an **even triple**. If $o(x), o(y)$ and $o(z)$ are all odd then we call (x, y, z) an **odd triple**. It is clear that since $z = (xy)^{-1}$, being generated by x, y and z is equivalent to being generated by x and y. If the context is clear we may refer to a hyperbolic generating triple simply as a 'triple' for brevity and write (x, y, xy) or even just (x, y).*

For a positive integer k let Q_{4k} be the dicyclic group of order $4k$ with presentation

$$Q_{4k} = \langle p, q \mid p^{2k} = q^4 = 1, p^q = p^{-1}, p^k = q^2\rangle.$$

Let $G = (H \times H) : Q_{4k}$ with the action of Q_{4k} defined as follows. For $(g_1, g_2) \in H \times H$ let $p(g_1, g_2) = (g_1, g_2)$ and $q(g_1, g_2) = (g_2, g_1)$. Then $G^0 = H \times H \times \langle p\rangle$ is a subgroup of index 2 inside G.

Theorem 9. *Let H be a perfect finite group. Let $a_1, c_1, a_2, c_2 \in H$ and, as before, define $\nu(a_i, c_i) = o(a_i)o(c_i)o(a_ic_i)$ for $i = 1, 2$. Assume that*

1. *the orders of a_1, c_1 are even,*
2. $\langle a_1, c_1\rangle = H$,
3. $\langle a_2, c_2\rangle = H$,
4. $\nu(a_1, c_1)$ *is coprime to* $\nu(a_2, c_2)$.

Set $k > 1$ to be any integer that divides $gcd(o(a_1), o(c_1)), G := (H \times H) : Q_{4k}$, $G^0 := H \times H \times \langle p\rangle$, $a{:=}(a_1, a_2, p)$ and $c := (c_1, c_2, p^{-1})$. Then $(G^0; a, c)$ is a mixed Beauville structure on G.

Proof. We verify that the conditions of Definition 1 are satisfied. Since k divides $\nu(a_1, c_1)$ it is coprime to $\nu(a_2, c_2)$ so we can clearly generate the elements $(1, a_2, 1)$ and $(1, c_2, 1)$ giving us the second factor. This also shows that we can generate the elements $a' = (a_1, 1, p)$ and $c' = (c_1, 1, p^{-1})$. Since H is perfect we can then generate the first factor. Finally, since we can generate $H \times H$ we can clearly generate $\langle p\rangle$, hence we satisfy condition $M1$.

Now let $g \in G \setminus G^0$ and $\gamma \in G^0$. Then $g\gamma$ is of the form $(h_1, h_2, q^i p^j)$ for some $h_1, h_2 \in H$, $i = 1, 3$ and $1 \leq j \leq 2k$. Then

$$(g\gamma)^2 = (h_1h_2, h_2h_1, (q^ip^j)^2) = (h_1h_2, h_2h_1, p^k).$$

For a contradiction, suppose that $(g\gamma)^2 \in \Sigma(a, c)$. Then since h_1h_2 has the same order as h_2h_1 condition 4 implies that $(g\gamma)^2 = (1, 1, p^k) \in \Sigma(a, c)$ if and only if k does not divide $o(a)$ or k does not divide $o(c)$. Note that if $(g\gamma)^2$ were a power of ac by construction it would be 1_G. Since by hypothesis k divides $\gcd(o(a_1), o(c_1))$ we satisfy conditions $M2$ and $M3$.

Finally, to show that condition $M4$ is satisfied, suppose $g' \in \Sigma(a, c) \cap \Sigma(a^g, c^g)$ for $g \in G \setminus G^0$. Since conjugation by such an element g interchanges the first two factors of any element, we again have from condition 4 that g' is of the form $(1, 1, p^i)$ for some power of p, but from our previous remarks it is clear that $p^i = 1_H$ and so $g' = 1_G$. □

Remark 10. In the proof of Theorem 9 we chose $a := (a_1, a_2, p)$ and $c := (c_1, c_2, p^{-1})$ but in principle we could have chosen the third factor of a or c to be 1 and the third factor in their product ac to be p or p^{-1} as appropriate. If we then required k to divide $\gcd(o(a), o(ac))$ or $\gcd(o(ac), o(c))$ as necessary this gives rise to further examples of mixed Beauville groups.

We conclude this section with a number of results regarding hyperbolic generating triples for $G \times G$. The number of prime divisors of the order of a group will pose an obvious restriction on finding a mixable Beauville structure and so naturally we would like to know for how large an n we can generate G^n with

a triple of a given type. Hall treats this question in much more generality in [22], here we simply mention that for a hyperbolic generating triple (x, y, z) of a given type this depends on how many orbits there are of triples of the same type under the action of $\mathrm{Aut}(G)$.

Definition 11. *Let G be a finite group and (a_1, b_1, c_1), (a_2, b_2, c_2) be hyperbolic generating triples for G. We call these two triples **equivalent** if there exists $g \in Aut(G)$ such that $\{a_1^g, b_1^g, c_1^g\} = \{a_2, b_2, c_2\}$.*

Since conjugate elements must have the same order the following is immediate.

Lemma 12. *Let G be a finite group and (a_1, b_1, c_1) a hyperbolic generating triple of G of type (l_1, m_1, n_1) and (a_2, b_2, c_2) a hyperbolic generating triple of G of type (l_2, m_2, n_2). If $\{l_1, m_1, n_1\} \neq \{l_2, m_2, n_2\}$ then (a_1, b_1, c_1) and (a_2, b_2, c_2) are inequivalent triples.*

Lemma 13. *Let G be a nonabelian finite group and let (a_1, b_1, c_1) and (a_2, b_2, c_2) be equivalent hyperbolic generating triples for G for some $g \in Aut(G)$. Then*

1. *if $a_1^g = a_2$ then $b_1^g = b_2$ and $c_1^g = c_2$,*
2. *if $a_1^g = b_2$ then $b_1^g = c_2$ and $c_1^g = a_2$, and;*
3. *if $a_1^g = c_2$ then $b_1^g = a_2$ and $c_1^g = b_2$.*

Proof. Note that in this instance we take $a_i b_i c_i = 1_G$ for $i = 1, 2$. Let $a_1^g = a_2$ and suppose $b_1^g = c_2 = (a_2 b_2)^{-1}$ and $c_1^g = b_2$. Then

$$b_2 = c_1^g = (b_1^{-1} a_1^{-1})^g = c_2^{-1} a_2^{-1} = a_2 b_2 a_2^{-1}$$

which implies that G is generated by elements that commute, a contradiction since G is nonabelian. The proof is analogous for the remaining two statements. □

Lemma 14. *Let G be a finite group, (x, y, z) a hyperbolic generating triple for G and $gcd(o(x), o(y)) = 1$. Then $((x, y), (y, x^y), (z, z))$ is a hyperbolic generating triple for $G \times G$.*

Proof. If (x, y, z) is a hyperbolic generating triple of G then so is (y, x^y, z) since $x^y = y^{-1}xy$. Then, since the orders of x and y are coprime we can can produce the elements $(x, 1_G), (y, 1_G), (1_G, y)$ and $(1_G, x^y)$ which generate $G \times G$. □

Remark 15. The proof of the preceding Lemma naturally generalises to any pair, or indeed triple, of elements in a hyperbolic generating triple whose orders are coprime.

3. The Alternating groups

We make heavy use of the following Theorem due to Jordan.

Theorem 16 (Jordan)**.** *Let G be a primitive permutation group of finite degree n, containing a cycle of prime length fixing at least three points. Then $G \geqslant A_n$.*

Lemma 17. *The alternating group A_6 and $A_6 \times A_6$ are mixable.*

Proof. For our even triples we take the following elements in the natural representation of A_6

$$x_1 = (1,2)(3,4,5,6),\ y_1 = (1,5,6,4)(2,3),\ \ x_1' = (1,2)(3,4,5,6),\ y_1' = (1,5,6).$$

It can easily be checked in GAP [19] that $(x_1, y_1, x_1 y_1)$ is an even triple for A_6 of type $(4, 4, 4)$ and that $((x_1, x_1'), (y_1, y_1'), (x_1 y_1, x_1' y_1'))$ is an even triple for $A_6 \times A_6$ of type $(4, 12, 12)$.

For our odd triples, let $x_2 = (1, 2, 3, 4, 5)$, $y_2 = x_2^{(1,3,6)}$ and $y_2' = x_2^{(1,2,3,4,6)}$. Then it can be checked that $(x_2, y_2, x_2 y_2)$ is an odd triple of type $(5, 5, 5)$ for A_6 and $((x_2, x_2), (y_2, y_2'), (x_2 y_2, x_2 y_2'))$ is an odd triple for $A_6 \times A_6$ also of type $(5, 5, 5)$. Therefore we have a mixable Beauville structure of type $(4, 4, 4; 5, 5, 5)$ for A_6 and type $(4, 12, 12; 5, 5, 5)$ for $A_6 \times A_6$. □

Lemma 18. *The alternating group A_7 and $A_7 \times A_7$ are mixable.*

Proof. For our even triples we take the following elements in the natural representation of A_7

$$x_1 = (1,2)(3,4)(5,6,7),\ y_1 = (1,2,3)(4,5)(6,7),\ \ x_1' = (1,6)(2,4,5)(3,7),\ y_1' = (1,6,2)(3,7,4).$$

It can easily be checked in GAP that $(x_1, y_1, x_1 y_1)$ is an even triple for A_7 of type $(6, 6, 5)$ and that $((x_1, x_1'), (y_1, y_1'), (x_1 y_1, x_1' y_1')$ is an even triple for $A_7 \times A_7$ of type $(6, 6, 5)$.

For our odd triples let $x_2 = (1, 2, 3, 4, 5, 6, 7)$, $y_2 = x_2^{(1,3,2)}$ and $y_2' = x_2^{(1,3,2)}$. Again, it can be checked that $(x_1, y_1, x_1 y_1)$ is an odd triple of type $(7, 7, 7)$ for A_7 and that $((x_1, x_1'), (y_1, y_1'), (x_1 y_1, x_1' y_1'))$ is an odd triple also of type $(7, 7, 7)$ for $A_7 \times A_7$. Therefore both A_7 and $A_7 \times A_7$ admit a mixable Beauville structure of type $(6, 6, 5; 7, 7, 7)$. □

Lemma 19. *The alternating group A_{2m} and $A_{2m} \times A_{2m}$ are mixable for $m \geq 4$.*

Proof. For $m \geq 4$ let $G = A_{2m}$ under its natural representation and consider the elements

$$\begin{aligned} a_1 &= (1,2)(3,\ldots,2m), \\ b_1 &= a_1^{(1,3,4)} = (1,5,6,\ldots,2m,4)(2,3), \\ a_1b_1 &= (1,3)(2,5,7,\ldots,2m-3,2m-1,4,6,8,\ldots,2m). \end{aligned}$$

The subgroup $H_1 = \langle a_1, b_1 \rangle$ is clearly transitive and the elements

$$a_1^2 = (3,5,\ldots,2m-1)(4,6,\ldots,2m), \;\; b_1^2 = (1,6,8,\ldots,2m)(5,7,\ldots,2m-1,4),$$

fix the point 2 and act transitively on the remaining points. Finally, $a_1^2 b_1^{-2} = (1,2m,2m-1,3,4)$ is a prime cycle fixing at least 3 points for all m and so by Jordan's Theorem $H_1 = G$. This gives us our first hyperbolic generating triple of type $(2m-2, 2m-2, 2m-2)$ for G. For our second, we show that there is a similar triple which is inequivalent to the first under the action of $\mathrm{Aut}(G) = S_{2m}$. Consider the elements

$$\begin{aligned} a_1' &= (1,2)(3,\ldots,2m), \\ b_1' &= a_1'^{(1,4,3)} = (1,3,5,6,\ldots,2m)(2,4), \\ a_1'b_1' &= (1,4,6,\ldots,2m,5,7,\ldots,2m-1)(2,3) \end{aligned}$$

and note that $a_1' = a_1$. For the same argument as before we have that $\langle a_1', b_1' \rangle = G$. Now suppose that (a_1, b_1, a_1b_1) is equivalent to $(a_1', b_1', a_1'b_1')$ for some $g \in \mathrm{Aut}(G)$. If $a_1^g = a_1'$ then by Lemma 13 we have that $b_1^g = b_1'$ and $(a_1b_1)^g = a_1'b_1'$. Then $(1,2)^g = (1,2)$ and $(2,3)^g = (2,4)$ but these are incompatible with $(1,3)^g = (2,3)$ since for the former to hold 3 must map to 4 which is incompatible with the latter. Now suppose $a_1^g = b_1'$ implying $b_1^g = a_1'b_1'$ and $a_1b_1 = a_1$. Then similarly we have $(1,2)^g = (2,4)$ and $(2,3)^g = (2,3)$ forcing g to map 1 to 4 which is incompatible with requiring that $(1,3)^g = (1,2)$. Finally, if $a_1^g = a_1'b_1'$, forcing $b_1^g = a_1'$ and $a_1b_1 = b_1'$, we get that $(1,2)^g = (2,3)$ and $(2,3)^g = (12)$ and we find this is incompatible with $(1,3)^g = (2,4)$. Hence these two hyperbolic generating triples are inequivalent under the action of the automorphism group of G and so $((a_1, a_1'), (b_1, b_1'), (a_1b_1, a_1'b_1'))$ is a hyperbolic generating triple for $G \times G$ of type $(2m-2, 2m-2, 2m-2)$.

For our first odd triple consider the elements

$$\begin{aligned} a_2 &= (1,2,\ldots,2m-1), \\ b_2 &= a_2^{(1,2m,3)} = (1,4,5,\ldots,2m-1,2m,2), \\ a_2b_2 &= (2,3,5,7,\ldots,2m-1,4,6,8,\ldots,2m-2,2m) \end{aligned}$$

and let $H_2 = \langle a_2, b_2 \rangle$. We clearly have transitivity and 2-transitivity, hence H_2 is primitive. Since $a_2 b_2^{-1} = (1,2m,2m-1,2,3)$ is a prime cycle fixing at least 3 points for all m, again we can apply Jordan's Theorem and we have that $H_2 = G$. Then (a_1, b_1, a_2, b_2) is a mixable Beauville structure on G of type

$$(2m-2, 2m-2, 2m-2; 2m-1, 2m-1, 2m-1).$$

For our second odd triple consider the elements

$$\begin{aligned} a_2' &= (1,2,\ldots,2m-1), \\ b_2' &= a_2'^{(1,3,2m)} = (2,2m,4,\ldots,2m-1,3), \\ a_2'b_2' &= (1,2m,4,6,\ldots,2m-2,3,5,\ldots,2m-1) \end{aligned}$$

and note that $a_2' = a_2$. It follows that $\langle a_2, b_2 \rangle = G$ from a similar argument as before and so it remains to show that $\langle (a_2, a_2'), (b_2, b_2') \rangle = G \times G$. Let $g \in \mathrm{Aut}(G)$ and suppose that $a_2^g = a_2'$. Then by Lemma 13 and inspection of the fixed points of these triples we have g fixes $2m$ and maps $3 \to 1$ and $1 \to 2$ which is incompatible with $a_2^g = a_2'$. Similarly, if $a_2^g = b_2'$ then $2m \to 1$, $3 \to 2$ and $1 \to 2m$ but from $b_2^g = a_2'b_2'$ g must then map $3 \to 5$, a contradiction. Finally, if $a_2^g = a_2'b_2'$ we get the mappings $2m \to 2$, $3 \to 2m$ and $1 \to 1$. But since $a_2^g = a_2'b_2'$, g must then also map $3 \to 4$, a final contradiction. Then (a_2, b_2, a_2b_2) and $(a_2', b_2', a_2'b_2')$ are inequivalent hyperbolic generating triples both of type $(2m-1, 2m-1, 2m-1)$ on G and so we get a mixable Beauville structure on $G \times G$ of type

$$(2m-2, 2m-2, 2m-2; 2m-1, 2m-1, 2m-1).$$

□

Lemma 20. *The alternating group A_{2m+1} and $A_{2m+1} \times A_{2m+1}$ are mixable for $m \geq 4$.*

Proof. For $m \geq 4$ let $G = A_{2m+1}$ under its natural representation and consider the elements

$$\begin{aligned} a_1 &= (1,2)(3,4)(5,\ldots,2m+1), \\ b_1 &= (1,2,\ldots,2m-3)(2m-2,2m-1)(2m,2m+1), \\ a_1b_1 &= (1,3,5,\ldots,2m-1,2m+1,6,8,\ldots,2m-4). \end{aligned}$$

The subgroup $G_1 = \langle a_1, b_1 \rangle$ is clearly transitive and the elements

$$\begin{aligned} b_1a_1^2b_1^{-1} &= (4,6,8,\ldots,2m,5,7,\ldots,2m-5,2m-1,2m+1), \\ a_1b_1^2a_1^{-1} &= (2,4,2m+1,6,8,\ldots,2m-4,1,3,\ldots,2m-5) \end{aligned}$$

both fix the point $2m-3$ and act transitively on the remaining points; hence G_1 acts primitively. Finally, the element $a_1b_1^{-1} = (2,2m-3,2m-1,2m+1,4)$ is a prime cycle fixing at least 3 points for all $m \geq 4$ and so by Jordan's Theorem $G_1 = G$. This gives our first hyperbolic generating triple of type $(2(2m-3), 2(2m-3), 2m-3)$ for G. For our second even triple, we manipulate the first in the following way. Let

$$\begin{aligned} a_1' &= (1,2m-4,\ldots,6,2m+1,2m-1,\ldots,3), \\ b_1' &= (1,2)(3,4)(5,\ldots,2m+1), \\ a_1'b_1' &= (1,2m-3,\ldots,2)(2m-2,2m-1)(2m,2m+1). \end{aligned}$$

Since $b_1' = a_1$ and $a_1'b_1' = b_1^{-1}$ it is clear that $\langle a_1', b_1' \rangle = G$. Note also that $a_1' = b_1^{-1}a_1^{-1}$. To see that (a_1, b_1, a_1b_1) and $(a_1', b_1', a_1'b_1')$ are inequivalent, suppose for a contradiction there exists $g \in \mathrm{Aut}(G)$ for which these triples are equivalent. Since conjugation preserves cycle type it must be that $(a_1b_1)^g = a_1'$ which, by Lemma 13, implies that $a_1^g = b_1'$ and $b_1^g = a_1'b_1'$. This give the equality

$$a_1b_1^{-1} = b_1'(a_1'b_1') = a_1^gb_1^g = (a_1b_1)^g = a_1' = b_1^{-1}a_1,$$

a contradiction since otherwise G would be abelian. Then, $((a_1, a_1'), (b_1, b_1'))$ is an even triple for $G \times G$ of type $(2(2m-3), 2(2m-3), 2(2m-3))$.

For our first odd triple consider the elements

$$\begin{aligned} a_2 &= (1,2,\ldots,2m+1), \\ b_2 &= a_2^{(1,2,3)} = (1,4,5,\ldots,2m,2m+1,2,3), \\ a_2b_2 &= (1,3,5,\ldots,2m-1,2m+1,4,6,\ldots,2m-2,2m,2). \end{aligned}$$

The subgroup $G_2 = \langle a_2, b_2 \rangle$ is clearly transitive while the elements $b_2^{-1}a_2^2 = (1,5,6,\ldots,2m+1)(3,4)$ and $a_2b_2^{-1} = (1,2m+1,3)$ fix the point 2 and act transitively on the remaining points. Hence G_2 is primitive, with a prime cycle fixing at least 3 points, then by Jordan's Theorem $G_2 = A_{2m+1}$. This gives us an odd triple of type $(2m+1, 2m+1, 2m+1)$ for G and so since we have $\gcd(2(2m-3), 2m+1) = 1$ it follows that (a_1, b_1, a_2, b_2) is a mixable Beauville structure for A_{2m+1} of type

$$(2(2m-3), 2(2m-3), 2m-3; 2m+1, 2m+1, 2m+1).$$

For our second odd triple consider the cycles

$$\begin{aligned} x_2 &= (1,2,\ldots,2m-1), \\ y_2 &= x_2^{(1,2m,2,2m+1,3)} = (1,4,5,\ldots,2m-1,2m,2m+1), \\ x_2y_2 &= (1,2,3,5,\ldots,2m-1,4,6,\ldots,2m,2m+1) \end{aligned}$$

and let $H_2 = \langle x_2, y_2 \rangle$. We clearly have transitivity while the elements $[x_2, y_2] = (1,2m,4,5,2)$ and $x_2[x_2, y_2] = (2,3,5,6,\ldots,2m-1,2m,4)$ show that H_2 acts transitively on the stabiliser of the point $2m+1$ and contains a prime cycle fixing at least 3 points for all m. Then by Jordan's Theorem $H_2 = G$ and this gives us a second odd triple of type $(2m-1, 2m-1, 2m-1)$. Since it is clear that $2(2m-3)$ is coprime to both $2m-1$ and $2m+1$ we then have a mixable Beauville structure on $G \times G$ of type

$$(2(2m-3), 2(2m-3), 2(2m-3); 4m^2-1, 4m^2-1, 4m^2-1).$$

□

4. The groups of Lie type

We make use of theorems due to Zsigmondy, generalising a theorem of Bang, and Gow which we include here for reference. Throughout this section $q = p^e$ will denote a prime power for a natural number $e \geq 1$.

Theorem 21 (Zsigmondy [41] or Bang [1], as appropriate)**.** *For any positive integer $a > 1$ and $n > 1$ there is a prime number that divides $a^n - 1$ and does not divide $a^k - 1$ for any positive integer $k < n$, with the following exceptions:*

(1) *$a = 2$ and $n = 6$; and*
(2) *$a + 1$ is a power of 2, and $n = 2$.*

We denote a prime with such a property $\Phi_n(a)$.

Remark 22. The case where $a = 2$, $n > 1$ and not equal to 6 was proven by Bang in [1]. The general case was proven by Zsigmondy in [41]. Hereafter we shall refer to this as Zsigmondy's Theorem. A more recent account of a proof is given by Lünburg in [29].

Definition 23. *Let G be a group of Lie type defined over a field of characteristic $p > 0$, prime. A **semisimple** element is one whose order is coprime to p. A semisimple element is **regular** if p does not divide the order of its centraliser in G.*

Theorem 24 (Gow [21])**.** *Let G be a finite simple group of Lie type of characteristic p, and let g be a non-identity semisimple element in G. Let L_1 and L_2 be any conjugacy classes of G consisting of regular semisimple elements. Then g is expressible as a product xy, where $x \in L_1$ and $y \in L_2$.*

Remark 25. A slight generalisation of this result to quasisimple groups appears in [14, Theorem 2.6].

4.1. **Projective Special Linear groups $PSL_2(q) \cong A_1(q)$.** The projective special linear groups $PSL_2(q)$ are defined over fields of order q and have order $q(q+1)(q-1)/k$ where $k = \gcd(2, q+1)$. Their maximal subgroups are listed in [18].

Lemma 26. *Let $G = PSL_2(7)$. Then G^n admits a mixable Beauville structure if and only if $n = 1, 2$.*

Proof. The maximal subgroups of G are known [11, p.2] and these are subgroups isomorphic to S_4 or point stabilisers in the natural representation of G on 8 points. Hyperbolic generating triples cannot have type $(3,3,3)$, since $\frac{1}{3} + \frac{1}{3} + \frac{1}{3} \not< 1$, and similarly for types $(2,2,2)$, $(2,2,4)$ or $(2,4,4)$. The number of hyperbolic generating triples of type $(7,7,7)$ can be computed using GAP, but since it is equal to the order of $\mathrm{Aut}(G)$ we see from [22] that there is no triple of type $(7,7,7)$ for G^n when $n > 1$. Triples of type $(4,4,4)$ exist and any such triple generates G since elements of order 4 are not contained in point stabilisers and inside a subgroup isomorphic to S_4 the product of three elements of order 4 cannot be equal to the identity. We can compute the number of such triples from the structure constants and since this is twice the order of $\mathrm{Aut}(G)$ we have that there exists a hyperbolic generating triple of type $(4,4,4)$ on G and on $G \times G$. We then see that this is the maximum number of direct copies of G for which there exists a mixable Beauville structure.

For our odd triple we then take a triple of type $(7,7,3)$ which can be shown to exist by computing their structure constants and are seen to generate G since if they were to belong to a maximal subgroup then the product of two elements of order 7 would again have order 7. This gives a mixable Beauville structure of type $(4,4,4;7,7,3)$ on G. Finally, we then have mixable Beauville structures on $G \times G$ of type $(4,4,4;7,7,21)$ by Lemma 12 or alternatively of type $(4,4,4;7,21,21)$ by Lemma 14. □

Lemma 27. *Let $G = PSL_2(8)$. Then $G \times G$ admits a mixable Beauville structure.*

Proof. It can easily be checked in GAP that for G there exists hyperbolic generating triples of types $(2,7,7)$, $(3,3,9)$ and $(3,9,9)$. The two odd triples are inequivalent by Lemma 12 and by Lemma 14 we have that there exists a mixable Beauville structure of type $(14,14,7;3,9,9)$ on $G \times G$. □

Lemma 28. *Let $G = PSL_2(9)$. Then both G and $G \times G$ admit a mixable Beauville structure.*

Proof. This follows directly from the exceptional isomorphism $PSL_2(9) \cong A_6$ and Lemma 17. □

We make use of the following Lemmas:

Lemma 29. *Let $G = PSL_2(q)$ for $q = 7, 8$ or $q \geq 11$. Let $k = gcd(2, q+1)$ and $\phi(n)$ be Euler's totient function. Then, under the action of $Aut(G) = P\Gamma L_2(q)$ the number of conjugacy classes of elements of order $\frac{q+1}{k}$ in G is $\phi(\frac{q+1}{k})/2e$.*

Proof. Elements of order $\frac{q+1}{k}$ are conjugate to their inverse so there are $\phi(\frac{q+1}{k})/2$ conjugacy classes of elements of order $\frac{q+1}{k}$ in $PSL_2(q)$. The only outer automorphisms of G come from the diagonal automorphisms and the field automorphisms, but since diagonal automorphisms do not fuse conjugacy classes of semisimple elements we examine the field automorphisms. These come from the action of the Frobenius automorphism on the elements of the field $\mathbb{F}_q$ sending each entry of the matrix to its p-th power. The only fixed points of this action are the elements of the prime subfield $\mathbb{F}_p$ and so, since the entries on the diagonal of the elements of order $\frac{q+1}{k}$ are not both contained in the prime subfield we have that the orbit under this action has length e, the order of the Frobenius automorphism. We then get e conjugacy classes of elements of order $\frac{q+1}{k}$ inside of G fusing under this action. Hence under the action of the full automorphism group there are $\phi(\frac{q+1}{k})/2e$ conjugacy classes of elements of order $\frac{q+1}{k}$. □

Lemma 30. *For a prime power, $q = p^e \geq 13$, $q \neq 27$, let $q^+ = \frac{q+1}{k}$ where $k = gcd(2, q+1)$. Then $\frac{\phi(q^+)}{2e} > 1$ where $\phi(n)$ is Euler's totient function.*

Proof. Let $S = \{p^i, q^+ - p^i \mid 0 \leq i \leq e-1\}$ be a set of $2e$ positive integers less than and coprime to q^+ and whose elements are distinct when $q \geq 13$. To this set we add k which will depend on q. When $p = 2$ we let $k = 7$ since for all $e > 3$, $7 \notin S$ and $\gcd(7, q^+) = 1$. When $p = 3$ we let $k = 11$, then for $e > 3$, $11 \notin S$ and $\gcd(11, q^+) = 1$. Now consider the cases $q \equiv \pm 1 \bmod 4$ for $p \neq 2, 3$. When $q \equiv 1 \bmod 4$, let $k = q^+ - 2$. Since $p \neq 2$ we have $k \notin S$ and since q^+ is odd when $q \equiv 1 \bmod 4$ we have $\gcd(k, q^+) = 1$. Finally, when $q \equiv 3 \bmod 4$ then e must be odd. When $e > 2$ then $k = \frac{p-1}{2} \notin S$ and is coprime to q^+. When $e = 1$, $q^+ = 2^i m$ where $i > 0$ and m is odd. Then $\phi(q^+) = \phi(2^i)\phi(m) = 2^{i-1}\phi(m) > 2$ since $p > 11$. This completes the proof. □

Lemma 31. *Let G be the projective special linear group $PSL_2(q)$ where $q \geq 7$. Then,*

1. *there is a mixable Beauville structure for $G \times G$, and;*
2. *when $p \neq 2$ there is also a mixable Beauville structure for G.*

Proof. In light of Lemmas 26–28 we can assume that $q \geq 11$. Define $q^+ = \frac{q+1}{k}$, where $k = \gcd(2, q+1)$, and similarly for q^-. Jones proves in [26] that hyperbolic generating triples of type (p, q^-, q^-) exist for G when $q \geq 11$ and since $\gcd(p, q^-) = 1$ we immediately have, by Lemma 14, a hyperbolic generating triple for $G \times G$. We proceed to show that there exists a hyperbolic generating triple (x, y, z) for G of type (q^+, q^+, q^+) and note that both p and q^- are coprime to q^+. The only maximal subgroups containing elements of order q^+ are the dihedral groups of order $2q^+$ which we denote by D_{q^+}. By Gow's Theorem, for a conjugacy class, C, of elements of order q^+ there exist $x, y, z \in C$ such that $xyz = 1$. Since inside D_{q^+} any conjugacy class of elements of order q^+ contains only two elements, x, y and z can not all be contained in the same maximal subgroup of G. Hence (x, y, z) is a hyperbolic generating triple for G of type (q^+, q^+, q^+). When the number of conjugacy classes of elements of order q^+ in G under the action of $\mathrm{Aut}(G)$ is strictly greater than 1 we can apply Gow's Theorem a second time to give a hyperbolic generating triple of type (q^+, q^+, q^+) for $G \times G$. This follows from Lemmas 29 and 30 with the exceptions of $q = 11$ or 27. For $G = PSL_2(11)$ we have that a triple of type (p, q^-, q^-) exists by [26] or alternatively the words ab and $[a, b]$ in the standard generators for G [40] give an odd triple of type $(11, 5, 5)$. In both cases we have, by Lemma 14, an odd triple of type or $(55, 55, 5)$ for $G \times G$. For our even triple, the structure constants for the number of triples of type $(6, 6, 6)$ can be computed and is seen to be twice the order of $\mathrm{Aut}(G)$ and so we have an even triple for G and $G \times G$. For $G = PSL_2(27)$ we take the words in the standard generators [40] $(ab)^2(abb)^2$, a^{b^2} which give an even triple of type $(2, 14, 7)$ and the words b^2, b^a which give an odd triple of type $(3, 3, 13)$. Again, by Lemma 14, these give a mixable Beauville structure on $G \times G$. Finally, we remark that when $q \equiv \pm 1 \bmod 4$ we have that q^- and q^+ have opposite parity and this determines the parity of our triples. When $q \equiv 1 \bmod 4$, (p, q^-, q^-) becomes our even triple, (q^+, q^+, q^+) our odd triple and vice versa when $q \equiv 3 \bmod 4$. □

4.2. **Projective Special Unitary groups $PSU_3(q) \cong {}^2A_2(q)$.** The projective special unitary groups $PSU_3(q)$ are defined over fields of order q^2 and have order $q^3(q^3+1)(q^2-1)/d$ where $d = (3, q+1)$. Their maximal subgroups can be found in [31] and we refer to the character table and notation in [34].

Lemma 32. *Let $G = PSU_3(q)$ for $q = 4$ or $q \geq 7$. Let $d = gcd(3, q+1)$ and $t' = \frac{q^2-q+1}{d}$. Then there exists a hyperbolic generating triple of type (t', t', t') for G.*

Proof. Let G, d and t' be as in the hypothesis. Let C be a conjugacy class of elements of order t' in G and for $g \in C$, let $T = \langle g \rangle$. When $q \geq 7$ the unique maximal subgroup of G containing g is $N_G(T)$ of order

$3t'$. Since $\gcd(6,t')=1$ and since T is by definition normal in $N_G(T)$ the Sylow p-subgroup in $N_G(T)$ for any prime $p|t'$ is contained in T and is therefore unique. Hence for all $x \in N_G(T) \setminus T$ the order of x is 3. Then for $g \in C$ since g is conjugate to g^{-q} and g^{q^2} [34] we have $C \cap N_G(T) = \{g, g^{-q}, g^{q^2}\}$. This partitions C into $|C|/3$ disjoint triples. Using the structure constants obtained from the character table of G we show that it is possible to find a hyperbolic generating triple of G entirely contained within C. Our method is to count the total number of triples $(x,y,z) \in C \times C \times C$ such that $xyz=1$, which we denote $n(C,C,C)$, and show that there exists at least one such triple where x,y,z come from distinct maximal subgroups. Let S be a triple of the form $\{g, g^{-q}, g^{q^2}\} \subset C$, then for $s_1, s_2, s_3 \in S, s_1s_2s_3 = 1$ if and only if all three elements are distinct. Therefore the contribution to $n(C,C,C)$ from triples contained within a single maximal subgroup of G is $2|C|$. Using the formula for structure constants as found in [16] we have that

$$n(C,C,C) = \frac{|C|^3}{|G|}\left(1 - \frac{1}{q(q-1)} - \frac{1}{q^3} - \sum_{1}^{t'-1} \frac{(\zeta_{t'}^{u} + \zeta_{t'}^{-uq} + \zeta_{t'}^{uq^2})^3}{3(q+1)^2(q-1)}\right)$$

where $\zeta_{t'}$ is a primitive t'-th root of unity. From the triangle inequality we have $|(\zeta_{t'}^{u} + \zeta_{t'}^{-uq} + \zeta_{t'}^{uq^2})|^3 \leq 27$ so we can bound $n(C,C,C)$ from below and for $q \geq 8$ the following inequality holds

$$n(C,C,C) \geq \frac{|C|^3}{|G|}\left(1 - \frac{1}{q(q-1)} - \frac{1}{q^3} - \frac{9(t'-1)}{(q+1)^2(q-1)}\right) > 2|C|.$$

For $q = 4$ or 7 direct computation of the structure constants show that we can indeed find a hyperbolic generating triple of the desired type. □

Lemma 33. *Let $G = PSU_3(q)$ for $q = 4$ or $q \geq 7$. Let $c = gcd(3, q^2-1)$, $d = gcd(3, q+1)$ and $t' = (q^2-q+1)/d$. Then*

(1) *for $p = 2$ there exists a mixable Beauville structure on G of type*

$$\left(2, 4, \frac{q^2-1}{c}; t', t', t'\right),$$

(2) *for $p \neq 2$ there exists a mixable Beauville structure on G of type*

$$\left(p, q+1, \frac{q^2-1}{d}; t', t', t'\right).$$

Proof. The existence of hyperbolic generating triples of type (t',t',t') in both even and odd characteristic is given in Lemma 32 and so we turn to the even triples. When $p=2$ we have the existence of our even triples from [14, Lemma 4.20 and Theorem 4.22] and so we now assume that $G = PSU_3(q)$ where p is odd, $q \geq 7$, $r = q+1$ and $s = q-1$.

From the list of maximal subgroups of G elements of order rs/d exist and can belong to subgroups corresponding to stabilisers of isotropic points, stabilisers of non-isotropic points and possibly one of the maximal subgroups of a fixed order which can occur is $PSU_3(q)$ for certain q. Stabilisers of isotropic points have order q^3r/d whereas stabilisers of non-isotropic points have order qr^2s/d. There exist $1+d$ conjugacy classes of elements of order p in G which are as follows. The unique conjugacy class, C_2, of elements whose centralisers have order q^3r/d; and d conjugacy classes, $C_3^{(l)}$ for $0 \leq l \leq d-1$, of elements whose centralisers have order q^2. Since an element of order p which stabilises a non-isotropic point belongs to a subgroup of G isomorphic to $SL_2(q)$, the order of its centraliser in G must be a multiple of $2p$, hence must belong to C_2. In particular, elements of C_3^0 do not belong to stabilisers of non-isotropic points. There exists a conjugacy class, C_6, of elements of order r whose centralisers have order r^2/d. An element of order r contained in the stabiliser of an isotropic point must be contained in a cyclic subgroup of order rs/d, hence rs/d must divide the order of its centraliser and so elements of C_6 are not contained in the stabilisers of isotropic points. Let C_7 be a conjugacy class of elements of order rs/d, then, any triple of elements $(x,y,z) \in C_3^0 \times C_6 \times C_7$, such that $xyz=1$, will not be entirely contained within the stabiliser of an isotropic or non-isotropic point. Let $n(C_3^0, C_6, C_7)$ be the number of such triples, then using the structure constant formula from [16] and the character table for G we have

$$n(C_3^0, C_6, C_7) = \frac{|C_3^0||C_6||C_7|}{|G|}\left(1 + \sum_{u=1}^{\frac{r}{d}-1} \frac{\epsilon^{3u}(\epsilon^{3u} + \epsilon^{6u} + \epsilon^{(r-3)u})}{t}\right)$$

where ϵ is a primitive r-th root of unity. Using the triangle inequality we can bound the absolute value of the summation by $\frac{3q}{q^2-q+1}$ which, for $q \geq 7$, is strictly less than 1. In order to show that such an (x,y,z)

is not contained in any of the possible maximal subgroups of order 36, 72, 168, 216, 360, 720 or 2520, notice that the subgroup generated by (x, y, z) has order divisible by $n = p(q^2 - 1)/d$, hence this can only occur when $p = 3$, 5 or 7. The only cases where $n \leq 2520$ and divides one of the possible subgroup orders are the cases $q = 7$ or 9, but none of these subgroups contain elements of order 48 or 80 so we see that this is indeed an even triple for G. Finally, we must show that $\gcd(\frac{4rs}{c}, t') = 1$ when p is even and $\gcd(\frac{prs}{d}, t') = 1$ when p is odd. For all p it is clear that $\gcd(p, t') = 1$ and so it suffices to show that $\gcd(rs, t') = 1$. We have that $t'd - s = q^2$ so t' is coprime to s and since $r^2 - t'd = 3q$ and t' is coprime to 3 we have that t' is coprime to r. □

In order to extend this to a mixable Beauville structure on $G \times G$ where $G = PSU_3(q)$ we will need the following Lemma:

Lemma 34. *Let $G = PSU_3(q)$ for $q \geq 3$ and $d = (3, q + 1)$. Then the number of conjugacy classes of elements of order $t' = (q^2 - q + 1)/d$ in G under the action of $Aut(G)$ is $\phi(t')/6e$ where $\phi(n)$ is Euler's totient function.*

Proof. Let $x \in G$ have order t', then x is conjugate to x^{-q} and x^{q^2} and so the number of conjugacy classes of order t' in G is $\phi(t')/3$. Then since the field automorphism has order $2e$ and the diagonal entries of an element of order t' are not all contained in the prime subfield its orbit has length $2e$. This gives the desired result. □

Lemma 35. *Let G be the projective special unitary group $PSU_3(q)$ for $q = 7$ or $q \geq 9$. Let $c = gcd(3, q^2 - 1)$, $d = gcd(3, q + 1)$ and $t' = (q^2 - q + 1)/d$. Then*

(1) *for $p = 2$ there exists a mixable Beauville structure on $G \times G$ of type*

$$\left(2\frac{q^2-1}{c}, 2\frac{q^2-1}{c}, 4; t', t', t'\right),$$

and;

(2) *for $p \neq 2$ there exists a mixable Beauville structure on $G \times G$ of type*

$$\left(p(q+1), p(q+1), p\frac{q^2-1}{d}; t', t', t'\right).$$

Proof. Let the conditions of the hypothesis be satisfied with $p = 2$. Then as in the proof of Lemma 33 there exists a hyperbolic generating triple of type $(2, 4, \frac{q^2-1}{c})$ on G and by Lemma 14 this yields an even triple of type $(2\frac{q^2-1}{c}, 2\frac{q^2-1}{c}, 4)$ on $G \times G$. Similarly, for $p \neq 2$ by Lemma 33 there exists a hyperbolic generating triple of type $(p, q + 1, \frac{q^2-1}{d})$ on G, which by Lemma 14 yields an even triple of type $(p(q + 1), p(q + 1), \frac{q^2-1}{d})$ on $G \times G$.

By Lemma 32 there exist hyperbolic generating triples of type (t', t', t') for all p and by Lemma 34 we need only show that $\phi(t') > 6e$. For $q = 7, 9$ it can be verified directly that $\phi(t') > 6e$ so we can assume $q \geq 11$. In the case $d = 1$ we have $2^3 < p^e - 1$ so $2^f < p^{2f} - p^f + 1$ for all $0 \leq f \leq 6e$ and we have our inequality. In the case $d = 3$ since $2^3 3 < p(p - 1)$ we have 2^f for $1 \leq f \leq 3e$. Similarly, since $3^2 < p - 1$ we have the terms 3^{f-1} for $1 \leq f \leq 4e$ giving us $7e$ terms in total, as was to be shown. □

Lemma 36. *The projective special unitary group $PSU_3(q)$ and $PSU_3(q) \times PSU_3(q)$ admit a mixable Beauville structure for $q \geq 3$.*

Proof. In light of the preceding Lemmas in this section it remains only to check the cases $PSU_3(3)$, $PSU_3(5)$ and $PSU_3(q) \times PSU_3(q)$ for $q = 3, 4, 5$ and 8. We present words in the standard generators [38, 40] that can be easily checked to give suitable triples for G which, by Lemma 14, give mixable Beauville structures for these cases. For $G = PSU_3(3)$ let

$$a_1 = [a, b^2],\ b_1 = [a, b^2]^b,\ a_2 = (babab^2)^3,\ b_2 = [a, b^2]^b \in G.$$

It can be checked that $G = \langle a_1, b_1\rangle = \langle a_2, b_2\rangle$ where both hyperbolic generating triples have type $(4, 4, 8)$ but in the former triple both elements of order 4 come from the conjugacy class $4C$, whereas in the latter, $a_2 \in 4AB, b_2 \in 4C$. Since these two triples are then inequivalent under the action of $\mathrm{Aut}(G)$ we have that $(a_1, a_2), (b_1, b_2) \in G \times G$ yields a hyperbolic generating triple of type $(4, 4, 8)$. For the remaining cases we present in Table 1 words in the standard generators for G which, by Lemma 14, give a mixable Beauville structure on $G \times G$ and G where necessary. □

q	x_1	y_1	x_2	y_2	Type
3	a_1, a_2	b_1, b_2	ab	ba	$(4, 4, 8; 7, 7, 3)$
4	a	$(ab)^2$	b	$[b, a]$	$(2, 13, 13; 3, 5, 3)$
5	a	ab^2	ab	b^3ab^3	$(3, 8, 8; 7, 7, 5)$
8	a	$(ab)^2$	$[a, b]$	$[a, b]^{babab}$	$(2, 19, 19; 9, 9, 7)$

TABLE 1. Words in the standard generators [40] a and b or as otherwise specified in Lemma 36 for $G = PSU_3(q)$.

4.3. **The Suzuki groups ${}^2B_2(2^{2n+1}) \cong Sz(2^{2n+1})$.** The Suzuki groups ${}^2B_2(q)$ are defined over fields of order $q = 2^{2n+1}$ for $n \geq 0$ and have order $q^2(q-1)(q^2+1)$. They are simple for $q > 2$ and their maximal subgroups can be found in [39].

Lemma 37. *Let G be the Suzuki group ${}^2B_2(q)$ for $q > 2$. Then*

(1) *G admits a mixable Beauville structure of type $(2, 4, 5; q-1, n, n)$, and;*
(2) *$G \times G$ admits a mixable Beauville structure of type $(4, 10, 10; n(q-1), n(q-1), n)$*

where $n = q \pm \sqrt{2q} + 1$, whichever is coprime to 5.

Proof. In the proof of [18, Theorem 6.2] Fuertes and Jones prove that there exist hyperbolic generating triples for G of types $(2, 4, 5)$ and $(q-1, n, n)$. It is clear that $\gcd(10, n) = \gcd(10, q-1) = 1$. Then, by Lemma 14, we need only show that $\gcd(q-1, n) = 1$. If $q-1$ and n share a common factor, then so do q^2-1 and q^2+1 and similarly their difference. Hence $\gcd(q-1, n)$ divides 2, but since $q-1$ is odd we have $\gcd(q-1, n) = 1$ as was to be shown. □

4.4. **The small Ree groups ${}^2G_2(3^{2n+1}) \cong R(3^{2n+1})$.** The small Ree groups are defined over fields of order $q = 3^{2n+1}$ for $n \geq 0$ and have order $q^3(q^3+1)(q-1)$. They are simple for $q > 3$ and their maximal subgroups can be found in [27] or [39].

Lemma 38. *The small Ree groups ${}^2G_2(q)$ for $q > 3$ admit a mixable Beauville structure of type*

$$\left(\frac{q+1}{2}, \frac{q+1}{2}, q + \sqrt{3q} + 1; \frac{q-1}{2}, \frac{q-1}{2}, q - \sqrt{3q} + 1\right).$$

Proof. Let $G = {}^2G_2(q)$ for $q > 3$ and for convenience we let $n^+ = q + \sqrt{3q} + 1$ and $n^- = q - \sqrt{3q} + 1$. Ward's analysis of the small Ree groups [36] shows that elements of orders $\frac{q+1}{2}$, $\frac{q-1}{2}$, n^+ and n^- exist and that the order of their centralisers are $q+1$, $q-1$, n^+ and n^-; hence these are all regular semisimple elements of G. Then, by Gow's Theorem, we can find elements $x_1, y_1 \in G$, both of order $\frac{q+1}{2}$, whose product has order n^+, and elements $x_2, y_2 \in G$, both of order $\frac{q-1}{2}$, whose product has order n^-.

The only maximal subgroups of G containing elements of order n^+ are normalisers of the cyclic subgroup which they generate of order $6n^+$. Similarly for elements of order n^-, they are contained in maximal subgroups of order $6n^-$. Since $n^+ - (q+1) = \sqrt{3q}$ we have $\gcd(\frac{q+1}{2}, n^+) = 1$ since neither is divisible by 3. Then, for $q > 3$ we have $\frac{q+1}{2} > 6$ so (x_1, y_1, x_1y_1) is indeed an even triple for G of type $(\frac{q+1}{2}, \frac{q+1}{2}, q + \sqrt{3q} + 1)$. Similarly, for $q > 3$ we have $\frac{q-1}{2} > 6$ and $n^+n^- + (q-1) = q^2$, hence $\gcd(\frac{q-1}{2}, n^-) = 1$ since again neither is divisible by 3. Note this also implies that $\gcd(n^+, \frac{q-1}{2}) = 1$. This gives us an odd triple for G of type $(\frac{q-1}{2}, \frac{q-1}{2}, q - \sqrt{3q} + 1)$.

It is clear that $\gcd(\frac{q+1}{2}, \frac{q-1}{2}) = 1$ since their difference is q and neither is divisible by 3. Similarly, $\gcd(n^+, n^-) = 1$ since their difference is $2\sqrt{3q}$ and both are clearly coprime to 6. Since we have already shown that $\gcd(n^+, \frac{q-1}{2}) = 1$ it remains to show that $\gcd(n^-, \frac{q+1}{2}) = 1$. We have $(q+1) - n^- = \sqrt{3q}$ and since neither is divisible by 3 we are done. □

Lemma 39. *The groups ${}^2G_2(q) \times {}^2G_2(q)$ for $q > 3$ admit a mixable Beauville structure of type*

$$\left(\frac{q+1}{2}, \frac{q+1}{2}n^+, \frac{q+1}{2}n^+; \frac{q-1}{2}, \frac{q-1}{2}n^-, \frac{q-1}{2}n^-\right)$$

where $n^+ = q + \sqrt{3q} + 1$ and $n^- = q - \sqrt{3q} + 1$.

Proof. By Lemmas 14 and 38 we need only show that $\gcd(\frac{q+1}{2}, n^+) = 1$ and $\gcd(\frac{q-1}{2}, n^-) = 1$, both of which were demonstrated in the proof of Lemma 38. □

4.5. **The large Ree groups ${}^2F_4(2^{2n+1})$.** The large Ree groups ${}^2F_4(q)$ are defined over fields of order $q = 2^{2n+1}$ for $n \geq 0$ and have order $q^{12}(q^6+1)(q^4-1)(q^3+1)(q-1)$. They are simple except for the case $q = 2$ which has simple derived subgroup ${}^2F_4(2)'$, known as the Tits group, which we consider along with the sporadic groups in the next section. The maximal subgroups of the large Ree groups can be found in [30] or [39].

Lemma 40. *The large Ree groups ${}^2F_4(q)$ for $q > 2$ admit a mixable Beauville structure of type*

$$\left(10, 10, n^+; \frac{q^2-1}{3}, n^-, n^-\right)$$

where $n^+ = q^2+q+1+\sqrt{2q}(q+1)$ and $n^- = q^2+q+1-\sqrt{2q}(q+1)$.

Proof. Let $G = {}^2F_4(q)$ for $q > 2$. Elements of order 10 exist since G contains maximal subgroups of the form ${}^2B_2(q) \wr 2$, as do elements of order $\frac{q^2-1}{3}$ since G contains maximal subgroups of the form $SU_3(q) : 2$ and $PGU_3(q) : 2$ and since $\gcd(3, q+1) = 3$. The only maximal subgroup containing an element of order n^+ is the normaliser of the cyclic subgroup which it generates, of order $12n^+$. Similarly, elements of order n^- are only contained in the normalisers of the cyclic subgroup they generate, of order $12n^-$.

Using the computer program CHEVIE it is possible to determine the structure constant for a pair of elements of order 10 whose product is n^+ and we see this is nonzero. Since $n^+n^- = q^4 - q^2 + 1$ and $q \equiv \pm 2 \bmod 5$ we have that $\gcd(10, n^-) = \gcd(10, n^+) = 1$. Then, since no maximal subgroup contains elements of orders 10 and n^+ this is indeed an even triple for G. Elements of order $\frac{q^2-1}{3}$ are semisimple and the order of the centraliser of an element of order n^- is n^- [33] so these are regular semisimple elements. Then by Gow's Theorem there exists a pair of elements of order n^- whose product has order $\frac{q^2-1}{3}$. To show that such a triple will generate G we show that no maximal subgroup contains elements of orders n^- and $\frac{q^2-1}{3}$. Since $(n^+n^-) + (q^2-1) = q^4$ any common factor of n^- and $\frac{q^2-1}{3}$ must be a power of 2, but since n^- and q^2-1 are both odd we have $\gcd(\frac{q^2-1}{3}, n^-) = 1$. Note that this also implies $\gcd(n^+, \frac{q^2-1}{3}) = 1$. Then, since $\frac{q^2-1}{3} > 12$ for $q > 2$ we then have that an odd triple of type $(\frac{q^2-1}{3}, n^-, n^-)$ exists for G.

We have already shown that $\gcd(10, n^-) = 1$, $\gcd(n^+, q^2-1) = 1$ and it is clear that $\gcd(10, \frac{q^2-1}{3}) = 1$. Finally, let $c = \gcd(n^+, n^-)$ and note that c is odd. Since $n^+ - n^- = 2\sqrt{2q}(q+1)$, c must divide $q+1$. Also, since $n^+ + n^- = 2(q^2+q+1)$, c must also divide q^2+q+1. Therefore c must divide q^2 and hence $c = 1$ so we have our desired mixable Beauville structure. □

Lemma 41. *The groups ${}^2F_4(q) \times {}^2F_4(q)$ for $q > 2$ admit a mixable Beauville structure of type*

$$\left(10, 10n^+, 10n^+; \frac{q^2-1}{3}n^-, \frac{q^2-1}{3}, n^-, n^-\right)$$

where $n^+ = q^2+q+1+\sqrt{2q}(q+1)$ and $n^- = q^2+q+1-\sqrt{2q}(q+1)$.

Proof. By Lemmas 14 and 40 we need only show that $\gcd(10, n^+) = 1$ and $\gcd(\frac{q^2-1}{3}, n^-) = 1$, both of which can be found in the proof of Lemma 40. □

4.6. **The Steinberg triality groups of type ${}^3D_4(q)$.** The Steinberg triality groups ${}^3D_4(q)$ are defined over fields of order q and have order $q^{12}(q^8+q^4+1)(q^6-1)(q^2-1)$. They are simple for all prime powers, q, and their maximal subgroups can be found in [28] or [39].

Lemma 42. *Let G be the Steinberg trialty group ${}^3D_4(2)$. Then both G and $G \times G$ admit a mixable Beauville structure.*

Proof. It can be verified using GAP that $(a, (ab)^3b^2; ab, b^{ab^2})$, where a and b are the standard generators as found in [40], is a mixable Beauville structure of type $(2, 7, 28; 13, 9, 13)$ and by Lemma 14 this yields a mixable Beauville strucure of type $(14, 14, 28; 117, 117, 13)$ on $G \times G$. □

Lemma 43. *Let G be the Steinberg triality group of type ${}^3D_4(q)$ for $q \geq 3$ and $d = gcd(3, q+1)$. Then*

(1) *for $p = 2$ there exists a mixable Beauville structure on G of type*

$$(6, 6, \Phi_{12}(q); \Phi_3(q), \Phi_3(q), \Phi_6(q)),$$

(2) *for $p \neq 2$ there exists a mixable Beauville structure on G of type*

$$\left(\frac{q^2-1}{d}, \frac{q^2-1}{d}, \Phi_{12}(q); \Phi_3(q), \Phi_3(q), \Phi_6(q)\right).$$

G	x_1	y_1	x_2	y_2
M_{11}	$ab(ab^2)^2$	$(ab(ab^2)^2)^b$	$(ab)^5$	$[a,b]^2$
M_{12}	$(ab)^4ba(bab)^2b$	$x_1^{b^2aba}$	ab	ba
J_1	$(ab)^2(ba)^3b^2ab^2$	b^{aba}	ab	$[a,b]$
M_{22}	$(ab)^3b^2ab^2$	$ba(bab^2)^2a$	ab	$(ab)^4b^2ab^2$
J_2	$ab(abab^2)^2ab^2$	$x_1^{ab^2}, x_1^{(ba)^2b^2}$	ab	ba
M_{23}	$abab^2$	$babab$	ab	$(abab)^{ba}$
${}^2F_4(2)'$	$(ab)^3bab$	$(x_1^7)^{baba}$	ab	ba
HS	$abab^3$	b^3aba	$(ab)^3b$	$((ab)^3b)^b$
J_3	$(ab)^2(ba)^3b^2$	x_1^b	ab	$(ab)^b$
M_{24}	$(ab)^4b$	$(x_1^3)^b$	ab	ba
McL	ab^2	bab	ab	$(ab)^b$
He	ab^3	ab^4	ab	ba
Ru	b	$b^{(ab)^5}$	$(ab)^2$	$(ba)^2$
Suz	$(ab(abab^2)^2)$	$x_1^{abab^2}$	ab	ba
$O'N$	$[a,b]$	$([a,b]^2)^{bab}$	ab^2	$(ab^2)^{abab}$
Co_3	ab	$(ab)^{b^2}$	$a(ab)^2b(a^2b)^2b$	$(x_2^3)^{baba}$
Co_2	a	b	$ab(ab^2)^2b$	$(x_2^2)^{bab^2}$
Fi_{22}	$(ab)^3b^3$	$((ab)^3b^3)^a$	b	b^a
HN	a	b	$[a,b]^a$	$(ab)^2b((ab)^5(ab^2)^2)^2$
Ly	a	b	$abab^3$	$x_2^{(ab)^7}$
Th	$[a,b]$	$[a,b]^{(ba)^4b^2}$	$ababa$	$(x_2^5)^{bab}$
Fi_{23}	a	b	$((ab)^{11}b)^3$	x_2^a
Co_1	a	b	$[(ab)^3, aba]$	$[(ab)^{23}, ab^2]^{babab}$
J_4	a	b	$a(bab)^3(ab)^2b$	x_2^a
Fi'_{24}	ab	$((ab)^6b)^{15}$	$b(ba)^3$	$(x_2^3)^{bab}$

TABLE 2. Words in the standard generators of G [40] where (x_1, y_1, x_2, y_2) is a mixable Beauville structure for G of the specified type.

Proof. Let G be as in the hypothesis. By [14, Lemma 5.24] for $q > 2$ there exists a hyperbolic generating triple of type $(\Phi_3(q), \Phi_3(q), \Phi_6(q))$.

For $p = 2$ one can verify using CHEVIE to compute the structure constants that there exist pairs of elements of order 6 whose product has order $\Phi_{12}(q)$ and it is clear from the list of maximal subgroups that this is indeed an even triple for G.

For $p \neq 2$ elements of order $\frac{q^2-1}{d}$ exist since G contains subgroups isomorphic to $SU(3,q)$. Using CHEVIE and the list of maximal subgroups it can be shown that an odd triple of type $(\frac{q^2-1}{d}, \frac{q^2-1}{d}, \Phi_{12}(q))$ exists for G. By Zsigmondy's theorem, Theorem 21, it is also clear that we have coprimeness for both Beauville structures. □

Lemma 44. *Let G be the Steinberg triality groups of type ${}^3D_4(q)$ for $q \geq 3$ and $d = gcd(3, q+1)$. Then*

(1) *for $p = 2$ there exists a mixable Beauville structure on $G \times G$ of type*

$$(6, 6\Phi_{12}(q), 6\Phi_{12}(q); \Phi_3(q), \Phi_3(q)\Phi_6(q), \Phi_3(q)\Phi_6(q));$$

(2) *for $p \neq 2$ there exists a mixable Beauville structure on G of type*

$$\left(\frac{q^2-1}{d}, \Phi_{12}(q)\frac{q^2-1}{d}, \Phi_{12}(q)\frac{q^2-1}{d}; \Phi_3(q), \Phi_3(q)\Phi_6(q), \Phi_3(q)\Phi_6(q)\right).$$

Proof. By Lemmas 14 and 43 we need only verify that $\gcd(6, \Phi_{12}(q)) = 1$ for $p = 2$ as the rest follows by construction. This is clear since $\Phi_{12}(q)$ is both odd and coprime to $q^2 - 1$ which is divisible by 3. □

5. The Sporadic groups

In this section we exhibit explicit words in the standard generators [40] of mixable Beauville structures for the sporadic groups. With the exception of the even triple for Janko's group, J_2, at least two elements of every triple have coprime order which, by Lemma 14, automatically gives us a corresponding triple for $G \times G$. In the case of J_2 we have two even triples, $(x_1, x_1^{ab^2})$ and $(x_1, x_1^{(ab)^2b^2})$ of types $(10, 10, 10)$

G	Type of G	Type of $G \times G$
M_{11}	$(8,8,5;11,3,11)$	$(8,40,40;11,33,33)$
M_{12}	$(8,8,5;11,11,3)$	$(8,40,40;11,33,33)$
J_1	$(10,3,10;7,19,19)$	$(10,30,30;133,133,19)$
M_{22}	$(8,8,5;11,7,7)$	$(8,40,40;77,77,7)$
J_2	$(10,10,10;7,7,3)$	$(10,10,80;7,21,21)$
M_{23}	$(8,8,11;23,23,7)$	$(8,88,88;23,161,161)$
${}^2F_4(2)'$	$(8,8,5;13,13,3)$	$(8,40,40;13,39,39)$
HS	$(8,8,15;7,7,11)$	$(8,120,120;7,77,77)$
J_3	$(8,8,5;19,19,3)$	$(8,40,40;19,57,57)$
M_{24}	$(8,8,5;23,23,3)$	$(8,40,40;23,69,69)$
McL	$(12,12,7;11,11,5)$	$(84,84,12;55,55,11)$
He	$(8,8,5;17,17,7)$	$(8,40,40;17,119,119)$
Ru	$(4,4,29;13,13,7)$	$(4,116,116;13,91,91)$
Suz	$(8,8,7;13,13,3)$	$(8,56,56;13,39,39)$
$O'N$	$(12,6,31;19,19,11)$	$(12,186,186;209,209,19)$
Co_3	$(14,14,5;23,23,9)$	$(14,70,70;23,207,207)$
Co_2	$(2,5,28;23,23,9)$	$(10,10,28;23,207,207)$
Fi_{22}	$(16,16,9;13,13,11)$	$(144,144,16;143,143,13)$
HN	$(2,3,22;5,19,19)$	$(6,6,22;95,95,19)$
Ly	$(2,5,14;67,67,37)$	$(10,10,14;2479,2479,67)$
Th	$(10,10,13;19,19,31)$	$(130,130,10;589,589,19)$
Fi_{23}	$(2,3,28;13,13,23)$	$(6,6,28;299,299,13)$
Co_1	$(2,3,40;11,13,23)$	$(6,6,40;143,143,23)$
J_4	$(2,4,37;43,43,23)$	$(4,74,74;989,989,43)$
Fi'_{24}	$(29,4,20;33,33,23)$	$(116,116,20;759,759,33)$
$\mathbb{B}$	$(2,3,8;47,47,55)$	$(6,6,8;47,2585,2585)$
$\mathbb{M}$	$(94,94,71;21,39,55)$	$(94,6674,6674;21,2145,2145)$

TABLE 3. Types of the mixable Beauville structures for G and $G \times G$ from the words in Table 2 and Lemmas 45 and 46.

and $(10,10,8)$, which are inequivalent by Lemma 12. For groups of reasonable order it can be explicitly computed in GAP that these elements generate. For groups of larger order we appeal to the list of maximal subgroups as found in [11] and [39] to ensure generation. For the Monster $\mathbb{M}$ and the Baby Monster $\mathbb{B}$ we use less constructive methods to prove the existence of these structures.

Lemma 45. *There exists a mixable Beauville structure on the Baby Monster, $\mathbb{B}$, and on $\mathbb{B} \times \mathbb{B}$.*

Proof. From [37] we know there exists a hyperbolic generating triple of type $(2,3,8)$. Let

$$x = (ab)^3(ba)^4b(ba)^2b^2, \;\; y = x^{ab^2}$$

be words in the standard generators [40]. They both have order 47 and their product has order 55, then from the list of maximal subgroups [39] they will generate $\mathbb{B}$. This gives a mixable Beauville structure of type $(2,3,8;47,47,55)$ on $\mathbb{B}$ and by Lemma 14 we have a mixable Beauville structure of type $(6,6,8;47,2585,2585)$ on $\mathbb{B} \times \mathbb{B}$. □

Lemma 46. *There exists a mixable Beauville structure on the Monster, $\mathbb{M}$, and on $\mathbb{M} \times \mathbb{M}$.*

Proof. Norton and Wilson [32, Theorem 21] show that the only maximal subgroups of the Monster which contain elements of order 94 are copies of $2 \cdot \mathbb{B}$ which does not contain elements of order 71. Then from a computation of the structure constants an even triple of type $(94,94,71)$ can be shown to exist. Finally, in [12] it is shown that there exists a hyperbolic generating triple of type $(21,39,55)$ on $\mathbb{M}$. Therefore we have a mixable Beauville structure of type $(94,94,71;21,39,55)$ on $\mathbb{M}$ and of type $(94,6674,6674;21,2145,2145)$. This completes the proof. □

Finally we have the following Lemma which completes the proof of Theorem 5.

Lemma 47. *Let G be one of the 26 sporadic groups or the Tits group ${}^2F_4(2)'$. Then there exists a mixable Beauville structure on G and $G \times G$.*

References

[1] A. S. Bang, 'Talteoretiske undersølgesler', *Tidskrift Math.* (5) 4 (1886) 130–137.

[2] N. Barker, N. Boston, N. Peyerimhoff and A. Vdovina, 'New examples of Beauville surfaces', *Monatsh. Math.* 166 (2012) 319–327.

[3] N. Barker, N. Boston, N. Peyerimhoff and A. Vdovina, 'An infinite family of 2–groups with mixed Beauville structures', *Int. Math. Res. Not.*, arXiv:math.AG/13044480.

[4] I. C. Bauer, 'Product–Quotient Suraces: Result and Problems', Preprint, 2012, arXiv:math.AG/12043409.

[5] I. C. Bauer, F. Catanese and F. Grunewald, 'Beauville surfaces without real structures I', *Geometric Methods in Algebra and Number Theory* Progr. Math. 235 (Birkhäuser Boston, Boston, 2005) 1–42.

[6] I. C. Bauer, F. Catanese and F. Grunewald, 'Chebycheff and Belyi polynomials, dessins d'enfants, Beauville surfaces and group theory', *Mediterr. J. Math.* (2) 3 (2006) 121–146.

[7] I. C. Bauer, F. Catanese and F. Grunewald, 'The classification of surfaces with $p_0 = q = 0$ isogenous to a product of curves', *Pure Appl. Math. Q.* 4 (2008) 547–586.

[8] I. C. Bauer, F. Catanese and R. Pignatelli, 'Surfaces of General Type with Geometric Genus Zero: A Survey', *Complex and differential geometry* Springer Proc. Math. 8 (Springer, Heidelberg, 2011) 1–48.

[9] I. C. Bauer, F. Catanese and R. Pignatelli, 'Complex surfaces of general type: some recent progress', *Global aspects of complex geometry* (Springer, Berlin, 2006) 1–58.

[10] F. Catanese, 'Fibered surfaces, varieties isogenous to a product and related moduli spaces', *Am. J. Math.* (1) 122 (2000) 1–44.

[11] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of Finite Groups* (Clarendon Press, Oxford, 1985).

[12] B. T. Fairbairn, 'Some Exceptional Beauville Structures', *J. Group Theory*, (5) 15 (2012) 631–639.

[13] B. T. Fairbairn, 'Recent work on Beauville surfaces, structures and groups', to appear in *Proceedings of Groups St Andrews 2013* (eds C.M. Campbell, M.R. Quick, E.F. Robertson, and C.M. Roney-Dougal).

[14] B. T. Fairbairn, K. Magaard and C. W. Parker, 'Generation of finite quasisimple groups with an application to groups acting on Beauville surfaces', *Proc. London Math. Soc.* (4) 107 (2013) 744–798.

[15] R. Friedman and J. W. Morgan 'Algebraic surfaces and four-manifolds: some conjectures and speculations', *Bull. Amer. Math. Soc.* (1) 18 (1988) 1–19.

[16] F. G. Frobenius, 'Über Gruppencharaktere', *Sitzungsber. Kön. Preuss. Akad. Wiss. Berlin* (1896) 985–1021.

[17] Y. Fuertes and G. González-Diez, 'On Beauville Structures on the Groups S_n and A_n', *Math. Z.* (4) 264 (2010) 959–968.

[18] Y. Fuertes and G. A. Jones, 'Beauville surfaces and finite groups', *J. Algebra* (1) 340 (2011) 13–27.

[19] The GAP Group, 'GAP – groups, algorithms and programming, version 4.7.4', 2014, http://www.gap-system.org.

[20] M. Geck, G. Hiss, F. Lübeck, G. Malle, and G. Pfeiffer, 'CHEVIE – A system for computing and processing generic character tables for finite groups of Lie type, Weyl groups and Hecke algebras', *Appl. Algebra Engrg. Comm. Comput.*, 7 (1996) 175–210.

[21] R. Gow, 'Commutators in finite simple groups of Lie type', *Bull. London Math. Soc.* (3) 32 (2000) 311–315.

[22] P. Hall, 'The Eulerian functions of a group', *Q. J. Math.* (1) 7 (1936) 134–151.

[23] G. A. Jones, 'Automorphism groups of Beauville surfaces', *J. Group Theory* (3) 16 (2013) 353–381.

[24] G. A. Jones, 'Beauville surfaces and groups: a survey', *Fields Inst. Comm.* to appear.

[25] G. A. Jones, 'Characteristically simple Beauville groups, I: cartesian powers of alternating groups', Preprint, 2013, arXiv:math.GR/13045444.

[26] G. A. Jones, 'Characteristically simple Beauville groups, II: low rank and sporadic groups', Preprint, 2013, arXiv:math.GR/13045450.

[27] P. B. Kleidman, 'The Maximal Subgroups of the Chevalley Groups $G_2(q)$ with q Odd, the Ree Groups ${}^2G_2(q)$, and Their Automorphism Groups', *J. Algebra* (1) 117 (1988) 30–71.

[28] P. B. Kleidman, 'The Maximal Subgroups of the Steinberg Triality Groups ${}^3D_4(q)$ and of Their Automorphism Groups', *J. Algebra* (1) 115 (1988) 182–199.

[29] H. Lünburg, 'Ein einfacher Beweis für den Satz von Zsigmondy über primitive Primteiler von $a^N - 1$', *Geometries and Groups* Lecture Notes in Mathematics 893 (eds Martin Aigner and Dieter Jungnickel; Springer, New York, 1981) 219–222.

[30] G. Malle, 'The maximal subgroups of ${}^2F_4(q^2)$', *J. Algebra* (1) 139 (1991) 52–69.

[31] H. Mitchell, 'Determination of the Ordinary and Modular Ternary Linear Groups', *Trans. Amer. Math. Soc.* (2) 12 (1911) 207–242.

[32] S. P. Norton and R. A. Wilson, 'Anatomy of the Monster. II', *Proc. London Math. Soc.* (3) 84 (2002) 581–598.

[33] K. Shinoda, 'The conjugacy classes of the finite Ree groups of type (F_4)', *J. Fac. Sci. Univ. Tokyo* Sect. 1 A, Mathematics, (1) 22 (1975) 1–15.

[34] W. A. Simpson and J. Sutherland Frame, 'The character tables for $SL(3,q)$, $SU(3,q^2)$, $PSL(3,q)$, $PSU(3,q^2)$', *Can. J. Math.* (3) XXV (1973) 486–494.

[35] J. Širáň, 'How symmetric can maps on surfaces be?', *Surveys in Combinatorics 2013* London Math. Soc. Lecture Note Ser. 409 (eds Simon R. Blackburn, Stefanie Gerke and Mark Wildon; CUP, Cambridge, 2013) 161–238.
[36] H. N. Ward 'On Ree's Series of Simple Groups' *Trans. Amer. Math. Soc.* (1) 121 (1966) 62–89.
[37] R. A. Wilson, 'The symmetric genus of the Baby Monster', *Q. J. Math.* (2) 44 (1993) 513–516.
[38] R. A. Wilson, 'Standard Generators for Sporadic Simple Groups', *J. Algebra* (2) 184 (1996) 505–515.
[39] R. A. Wilson, *The finite simple groups* Graduate Texts in Mathematics 251 (Springer-Verlag London Ltd., London, 2009).
[40] R. A. Wilson, *et al.*, Atlas of Finite Group Representations v3, http://brauer.maths.qmul.ac.uk/Atlas/v3.
[41] K. Zsigmondy, 'Zur Theorie der Potenzreste', *Monat. Math. Phys.* (1) 3 (1892) 265–284.

Department of Economics, Mathematics and Statistics, Birkbeck, University of London, Malet St, London WC1E 7HX, b.fairbairn@bbk.ac.uk, e.pierro@mail.bbk.ac.uk